Undoubtedly allied to Paxillus panuoides, but at once distin-

guished by the peculiar spores.

99. Panus strigosus, B. & C. Maximus; pileo excentrico depresso stipiteque valido strigosis; lamellis latis distantibus decurrentibus. Curt. nos. 6197, 5170. On oak stumps, New England, C. J. Sprague; Pennsylvania, Dr. Michener.

White; pileus 8 inches across, excentric, clothed with coarse strigose pubescence; margin thin; stem 2-3 inches high, 1 inch or more thick, strigose like the pileus; gills broad, distant,

decurrent.

Allied to Panus levis.

100. Panus operculatus, B. & C. Fasciculato-erumpens; pileo cupulæformi apice affixo rufo furfuraceo-villoso glabrescente; lamellis angustis tenuibus primum velo operculatis. Curt.

no. 5697. On bark, New England, D. Murray.

Fasciculate, erumpent; pileus cup-shaped, ½ an inch or more across, fixed by the apex, rufous, clothed with scurfy pubescence, which at length vanishes; gills narrow, of the same colour as the pileus, at first covered by a tympanoid veil.

Allied to P. Delastrii, Mont.

XXXII.—Notice of a new Antelope (Kobus) from Central Africa. By John Edward Gray, F.R.S.

THE British Museum has lately received, through Mr. Samuel Stevens, a number of heads and horns of Antelopes, &c., from Awan Bahr il Gazal (lat. 6° 70′ N.), in Central Africa, which have been brought to England by Mr. Petherick, who has been in the country collecting ivory, &c. I will first give a list of the animals brought, to show their geographical range, and then proceed to give a short notice of the new species.

1. Bos Taurus, Gray, Cat. of Ungulata in Coll. B. M. p. 17. Variety with the horns elongate and subspiral. Nader-Sobat, lat. 11° N.

2. Bubalus Caffer, Gray, Cat. p. 28. Horns.

- 3. Adenota Lechèe, Gray, Cat. p. 98. Male skin, in a bad state; skull of half-grown male and female. Awan and Raik.
- Kobus Maria, n. s. Two heads, male and female. Awan.
 Kobus ellipsiprymnus, p. 99. Head of male and female.

 Raik.
- 6. Alcelaphus Bubalis, p. 123. Several heads, of both sexes. Djour.

7. Damalis Senegalensis, p. 126. Head of male and females. Azack.

8. Tragelaphus sylvatica, p. 139. Head of male.

9. Giraffa Camelopardalis, p. 181. Head of male; pale variety.

10. Rhinoceros bicornis. Skull, with horns of a large size;

half-grown.

11. Hippopotamus amphibius. Skull. From Senaar, lat. 12° 13′ N.

I propose to name the new species, after my wife, who has so assisted me in my studies,



Head of male blackish-brown; lips, chin, gullet, orbits and temple enclosing the base of the ears, and the ears, whitish; the sides of the nose brownish; the hair of the cheek, side of the lower jaw, gullet, and upper part (all that remains) of the neck elongate, rigid; the horns elongate, rather slender, widely lyrate, with very strong transverse ridges and incurved tips.

Head of female brown, with the chin and gullet, base of the ears, and two small spots (one over the front and the other behind the lower edge of the eye) whitish; the hair of the head black, with brown tips; of the lower part of the cheek, lower

jaw, and neck very long. Hornless.